

Lower bound theorems for general polytopes

Guillermo Pineda, Julien Ugon & David Yost

SPCOM 2015

A polytope is the convex hull of a finite set.

They are encountered as the feasible regions for linear programming problems. The simplex algorithm works by moving from one vertex to another, so it is interesting to bound the lengths of paths in polytopes.

A polytope is the convex hull of a finite set.

They are encountered as the feasible regions for linear programming problems. The simplex algorithm works by moving from one vertex to another, so it is interesting to bound the lengths of paths in polytopes.

This motivated the Hirsch conjecture (1957): that any two vertices of a d -dimensional polytope with n facets must be connected by a path of length at most $n - d$. The conjecture is now known to be false in dimension 20. The best known upper bounds on the diameter are only sub-exponential in n and d .

A polytope is the convex hull of a finite set.

They are encountered as the feasible regions for linear programming problems. The simplex algorithm works by moving from one vertex to another, so it is interesting to bound the lengths of paths in polytopes.

This motivated the Hirsch conjecture (1957): that any two vertices of a d -dimensional polytope with n facets must be connected by a path of length at most $n - d$. The conjecture is now known to be false in dimension 20. The best known upper bounds on the diameter are only sub-exponential in n and d .

Here we attempt to estimate the total number of edges, in terms of the number of vertices.

A polytope is the convex hull of a finite set.

They are encountered as the feasible regions for linear programming problems. The simplex algorithm works by moving from one vertex to another, so it is interesting to bound the lengths of paths in polytopes.

This motivated the Hirsch conjecture (1957): that any two vertices of a d -dimensional polytope with n facets must be connected by a path of length at most $n - d$. The conjecture is now known to be false in dimension 20. The best known upper bounds on the diameter are only sub-exponential in n and d .

Here we attempt to estimate the total number of edges, in terms of the number of vertices.

It is also interesting to bound the total number of k -dimensional faces, but we will generally ignore the case $k > 1$ in this short presentation.

Precise upper bounds for the numbers of edges are easy to obtain. If $d = 3$, a polyhedron with v vertices has at most $3v - 6$ edges, with equality iff every face is a triangle. Such maximal examples are easy to construct.

Precise upper bounds for the numbers of edges are easy to obtain. If $d = 3$, a polyhedron with v vertices has at most $3v - 6$ edges, with equality iff every face is a triangle. Such maximal examples are easy to construct.

If $d \geq 4$, the cyclic polytope $C(v, d)$ has precisely $\binom{v}{2}$ edges. Clearly this is the maximum possible.

Precise upper bounds for the numbers of edges are easy to obtain. If $d = 3$, a polyhedron with v vertices has at most $3v - 6$ edges, with equality iff every face is a triangle. Such maximal examples are easy to construct.

If $d \geq 4$, the cyclic polytope $C(v, d)$ has precisely $\binom{v}{2}$ edges. Clearly this is the maximum possible.

McMullen (1970) established the corresponding conclusion for k -dimensional faces for all k ; this is known as the Upper Bound Theorem.

Lower bounds are not so easy to obtain. The following result of Barnette (1973) was considered a major breakthrough at the time. A polytope is *simplicial* if every facet (maximal proper face) is a simplex.

Theorem

A d -dimensional simplicial polytope v vertices has at least $dv - \binom{d}{2}$ edges; and there exist simplicial polytopes, namely the stacked polytopes, with precisely this many edges.

Lower bounds are not so easy to obtain. The following result of Barnette (1973) was considered a major breakthrough at the time. A polytope is *simplicial* if every facet (maximal proper face) is a simplex.

Theorem

A d -dimensional simplicial polytope v vertices has at least $dv - \binom{d}{2}$ edges; and there exist simplicial polytopes, namely the stacked polytopes, with precisely this many edges.

Barnette actually established the corresponding conclusion for k -dimensional faces for all k .

Lower bounds are not so easy to obtain. The following result of Barnette (1973) was considered a major breakthrough at the time. A polytope is *simplicial* if every facet (maximal proper face) is a simplex.

Theorem

A d -dimensional simplicial polytope v vertices has at least $dv - \binom{d}{2}$ edges; and there exist simplicial polytopes, namely the stacked polytopes, with precisely this many edges.

Barnette actually established the corresponding conclusion for k -dimensional faces for all k .

There are some estimates for cubical polytopes, but little seems to be known for general polytopes.

We denote by $\phi(v, d)$ the minimum possible number of edges, over all d -polytopes with v vertices.

It is well known that $\phi(v, 3)$ is either $3v/2$ or $\frac{1}{2}(3v + 1)$ depending on the parity of v (Steinitz, 1906). Examples achieving these bounds are easily constructed by successively slicing corners off a tetrahedron or a pyramid.

The 4-dimensional case was solved by Grünbaum in 1967. He showed that $\phi(6, 4) = 13$, $\phi(7, 4) = 15$, $\phi(10, 4) = 21$, and that $\phi(v, 4) = 2v$ for all other values of v .

The 4-dimensional case was solved by Grünbaum in 1967. He showed that $\phi(6, 4) = 13$, $\phi(7, 4) = 15$, $\phi(10, 4) = 21$, and that $\phi(v, 4) = 2v$ for all other values of v .

Note that a 4-dimensional simplex has 5 vertices and 10 edges and so $\phi(5, 4) = 10$. It is also *simple* in the sense that every vertex has degree 4; all such 4-polytopes have $\phi(v, 4) = 2v$.

The 4-dimensional case was solved by Grünbaum in 1967. He showed that $\phi(6, 4) = 13$, $\phi(7, 4) = 15$, $\phi(10, 4) = 21$, and that $\phi(v, 4) = 2v$ for all other values of v .

Note that a 4-dimensional simplex has 5 vertices and 10 edges and so $\phi(5, 4) = 10$. It is also *simple* in the sense that every vertex has degree 4; all such 4-polytopes have $\phi(v, 4) = 2v$.

Slicing a corner from a simple 4-polytope gives another simple polytope with three more vertices and six more edges. Thus we obtain simple polytopes with $v = 5, 8, 11, \dots$.

The 4-dimensional case was solved by Grünbaum in 1967. He showed that $\phi(6, 4) = 13$, $\phi(7, 4) = 15$, $\phi(10, 4) = 21$, and that $\phi(v, 4) = 2v$ for all other values of v .

Note that a 4-dimensional simplex has 5 vertices and 10 edges and so $\phi(5, 4) = 10$. It is also *simple* in the sense that every vertex has degree 4; all such 4-polytopes have $\phi(v, 4) = 2v$.

Slicing a corner from a simple 4-polytope gives another simple polytope with three more vertices and six more edges. Thus we obtain simple polytopes with $v = 5, 8, 11, \dots$.

Slicing an edge from a simple 4-polytope gives another simple polytope with four more vertices and eight more edges. Thus we obtain simple polytopes with $v = 9, 13, \dots, 12, 16, \dots$.

Simple polytopes in higher dimensions

A d -dimensional polytope is *simple* if every vertex has degree d .

For any polytope, the sum of the degrees of the vertices is equal to twice the number of edges. So in general $\phi(v, d) \geq \frac{1}{2}dv$, with equality only if there exists a simple polytope with v vertices. For which values of v do we find simple polytopes?

Simple polytopes in higher dimensions

A d -dimensional polytope is *simple* if every vertex has degree d .

For any polytope, the sum of the degrees of the vertices is equal to twice the number of edges. So in general $\phi(v, d) \geq \frac{1}{2}dv$, with equality only if there exists a simple polytope with v vertices. For which values of v do we find simple polytopes?

For $1 \leq k < d$, slicing off a $(k - 1)$ -dimensional face from a d -dimensional simple polytope will give another simple polytope with $kd - k^2$ more vertices. If d is even, then $d - 1$ and $2d - 4$ are relatively prime. Hence the following observation.

Theorem

If d is even, there is an integer K such that, for all $v > K$, $\phi(v, d) = \frac{1}{2}dv$ (i.e. there is a simple d -polytope with v vertices).

Theorem

If d is even, there is an integer K such that, for all $v > K$, $\phi(v, d) = \frac{1}{2}dv$ (i.e. there is a simple d -polytope with v vertices).

Even if d is odd, $d - 1$ and $2d - 4$ have no odd common prime factors.

Theorem

If d is odd, there is an integer K such that, for all even $v > K$, $\phi(v, d) = \frac{1}{2}dv$ (i.e. there is a simple d -polytope with v vertices).

Theorem

If d is even, there is an integer K such that, for all $v > K$, $\phi(v, d) = \frac{1}{2}dv$ (i.e. there is a simple d -polytope with v vertices).

Even if d is odd, $d - 1$ and $2d - 4$ have no odd common prime factors.

Theorem

If d is odd, there is an integer K such that, for all even $v > K$, $\phi(v, d) = \frac{1}{2}dv$ (i.e. there is a simple d -polytope with v vertices). Also, for all odd $v > K$, we have $\phi(v, d) = \frac{1}{2}d(v + 1) - 1$.

So the problem of calculating $\phi(v, d)$ is more interesting for small values of v .

A simplex shows that $\phi(d + 1, d) = \binom{d+1}{2}$. A prism based on a $(d - 1)$ -dimensional simplex shows that $\phi(2d, d) = d^2$.

A simplex shows that $\phi(d+1, d) = \binom{d+1}{2}$. A prism based on a $(d-1)$ -dimensional simplex shows that $\phi(2d, d) = d^2$.

Grünbaum proved that

$$\phi(d+k, d) = \binom{d}{2} - \binom{k}{2} + kd$$

for $1 \leq k \leq 4$,

A simplex shows that $\phi(d + 1, d) = \binom{d+1}{2}$. A prism based on a $(d - 1)$ -dimensional simplex shows that $\phi(2d, d) = d^2$.

Grünbaum proved that

$$\phi(d + k, d) = \binom{d}{2} - \binom{k}{2} + kd$$

for $1 \leq k \leq 4$, and conjectured that it holds in fact for every $k \leq d$.

A simplex shows that $\phi(d + 1, d) = \binom{d+1}{2}$. A prism based on a $(d - 1)$ -dimensional simplex shows that $\phi(2d, d) = d^2$.

Grünbaum proved that

$$\phi(d + k, d) = \binom{d}{2} - \binom{k}{2} + kd$$

for $1 \leq k \leq 4$, and conjectured that it holds in fact for every $k \leq d$.

We show that this is true, and moreover that the minimising polytope is unique.

A simplex shows that $\phi(d + 1, d) = \binom{d+1}{2}$. A prism based on a $(d - 1)$ -dimensional simplex shows that $\phi(2d, d) = d^2$.

Grünbaum proved that

$$\phi(d + k, d) = \binom{d}{2} - \binom{k}{2} + kd$$

for $1 \leq k \leq 4$, and conjectured that it holds in fact for every $k \leq d$.

We show that this is true, and moreover that the minimising polytope is unique.

We also obtain precise values for $\phi(2d + 1, d)$ and $\phi(2d + 2, d)$.

Theorem

Let P be a d -dimensional polytope with $d + k$ vertices, where $0 < k \leq d$.

(i) If P is $(d - k)$ -fold pyramid over the k -dimensional prism based on a simplex, then P has $\binom{d}{2} - \binom{k}{2} + kd$ edges.

(ii) Otherwise the numbers of edges is $> \binom{d}{2} - \binom{k}{2} + kd$.

Slicing one corner from the base of a square pyramid yields a polyhedron with 7 vertices and 6 faces, one of them a pentagon. We call this a *pentasm*.

We will use the same name for the higher-dimensional version, obtained by slicing one corner from the quadrilateral base of a $(d - 2)$ -fold pyramid. It has $2d + 1$ vertices and can also be represented as the Minkowski sum of a d -dimensional simplex, and a line segment which lies in the affine span of one 2-face but is not parallel to any edge.

Theorem

Let P be a d -dimensional polytope with $2d + 1$ vertices.

- (i) If P is d -dimensional pentasm, then P has $d^2 + d - 1$ edges.
- (ii) Otherwise the numbers of edges is $> d^2 + d - 1$, or P is the sum of two triangles.

This shows that the pentasm is the unique minimiser if $d \geq 5$.

If $d = 4$, the sum of two triangles has 9 vertices, and is the unique minimiser, with only 18 edges.

If $d = 3$, the sum of two triangles can have 7, 8 or 9 vertices; the example with $v = 7$ has 11 edges, the same as the pentasm.

Summarising, $\phi(9, 4) = 18$, and $\phi(2d + 1, d) = d^2 + d - 1$ for all $d \neq 4$.

Slicing one corner from the apex of a square pyramid yields a polyhedron combinatorially equivalent to the cube. Slicing one corner from 3-prism yields a polyhedron combinatorially equivalent to the 5-wedge. Of all the polyhedra with 8 vertices, these are the only two with 12 edges.

Slicing one corner from the apex of a square pyramid yields a polyhedron combinatorially equivalent to the cube. Slicing one corner from a 3-prism yields a polyhedron combinatorially equivalent to the 5-wedge. Of all the polyhedra with 8 vertices, these are the only two with 12 edges.

We show that for $d \neq 5$, analogues of these polyhedra minimise the number of edges, amongst polytopes with $2d + 2$ vertices.

Consider first the polytope obtained by slicing one corner from the apex of a $(d - 2)$ -fold pyramid on a square base. It has $2d + 2$ vertices, $(d + 1)^2 - 4$ edges and can also be represented as the Minkowski sum of a $(d - 3)$ -fold pyramid on a square base, and a line segment in the other dimension.

Slicing one corner from the apex of a square pyramid yields a polyhedron combinatorially equivalent to the cube. Slicing one corner from 3-prism yields a polyhedron combinatorially equivalent to the 5-wedge. Of all the polyhedra with 8 vertices, these are the only two with 12 edges.

We show that for $d \neq 5$, analogues of these polyhedra minimise the number of edges, amongst polytopes with $2d + 2$ vertices.

Consider first the polytope obtained by slicing one corner from the apex of a $(d - 2)$ -fold pyramid on a square base. It has $2d + 2$ vertices, $(d + 1)^2 - 4$ edges and can also be represented as the Minkowski sum of a $(d - 3)$ -fold pyramid on a square base, and a line segment in the other dimension.

Consider next a $(d - 3)$ -fold pyramid whose base is a 3-prism, then slice one corner off. This example also has $2d + 2$ vertices and $(d + 1)^2 - 4$ edges.

Theorem

Let P be a d -dimensional polytope with $2d + 2$ vertices, where $d \geq 6$ or $d = 3$.

(i) If P is one of the two polytopes just described, then P has $d^2 + 2d - 3$ edges.

(ii) Otherwise the numbers of edges is $> d^2 + 2d - 3$.

Theorem

Let P be a d -dimensional polytope with $2d + 2$ vertices, where $d \geq 6$ or $d = 3$.

(i) If P is one of the two polytopes just described, then P has $d^2 + 2d - 3$ edges.

(ii) Otherwise the numbers of edges is $> d^2 + 2d - 3$.

If $d = 4$, there is a third minimising polytope with 10 vertices and 21 edges.

If $d = 5$, the unique minimiser is the sum of a tetrahedron and triangle; this clearly has 12 vertices and 30 edges; $30 < 32$.

Summarising, $\phi(12, 5) = 30$, and $\phi(2d + 2, d) = d^2 + 2d - 3$ for all $d \neq 5$.

The case of $2d + 3$ vertices appears to be difficult.

Theorem

If $0 \leq k < d$, then

$$d^2 + \frac{1}{2}kd \leq \phi(2d + k, d) \leq d^2 + kd - \binom{k+1}{2}.$$

*The upper bound is the exact value if $k = 1, 2$ (unless $d = 4$ or 5).
The lower bound is the exact value if $k = 0, d - 3$. Being equal,
both are correct if $k = d - 1$.*

It is well known that there is no polyhedron with 7 edges. More generally a d -polytope cannot have between $\frac{1}{2}(d^2 + d - 2)$ and $\frac{1}{2}(d^2 + 3d - 4)$ vertices, inclusive. Grünbaum [p 188] discusses gaps in the possible number of edges, pointing that a second gap opens when $d = 6$ and a third gap opens when $d = 11$. Our main theorem shows that there are infinitely many gaps.

More precisely, in dimension $n^2 + 2$, there is no polytope with $\frac{1}{2}(n^4 + 2n^3 + 4n^2 + 3n + 4)$ edges.

The cyclic polytope $C(n^2 + n + 2, n^2 + 2)$ has one edge less, and the free join of an $(n^2 - n)$ -dimensional simplex and an $(n + 1)$ -prism has one edge more.

For example: a 27-dimensional polytope cannot have 497 edges. But there is a cyclic polytope with 496 edges, and a multiplex with 498 edges.

Finally, instead of just asking for upper and lower bounds, one may ask for the complete range of values. So for fixed d , can we describe for exactly which values of e, v there exists a d -polytope with v vertices and e edges?

If $d = 3$, the answer is well known: if and only if $\frac{3}{2}v \leq e \leq 3v - 6$.

For $d = 4$, the complete answer was given by Grünbaum: iff $2v \leq e \leq \binom{v}{2}$ **and** (v, e) is not one of the pairs $(6, 12)$, $(7, 14)$, $(10, 20)$ or $(8, 17)$.

The first three exceptions are clear: the number of vertices of a simple d -polytope cannot be between d and $2d$, and it can be $2d + 2$ only if $d = 5$.

The fourth case seemed like an oddity, but it is part of a general pattern. Here is a small part of that pattern.

Theorem

If there is a d -polytope with $2d$ vertices and $d^2 + 1$ edges, then $d = 3$.

More generally, we have the following. Define the *excess degree* of a polytope as

$$2e - dv = \sum_{v \in V} (\deg v - d).$$

Obviously a polytope is simple iff its excess degree is 0.

The fourth case seemed like an oddity, but it is part of a general pattern. Here is a small part of that pattern.

Theorem

If there is a d -polytope with $2d$ vertices and $d^2 + 1$ edges, then $d = 3$.

More generally, we have the following. Define the *excess degree* of a polytope as

$$2e - dv = \sum_{v \in V} (\deg v - d).$$

Obviously a polytope is simple iff its excess degree is 0.

Theorem

If P is a non-simple d -polytope, then its excess degree is at least $d - 2$.

For $d = 5$ or 6 , this helps us describe precisely the values of e, v for which there exists a d -polytope with v vertices and e edges.

Theorem

There is a simple 5-polytope with v vertices and e edges, iff $2e = 5v$ (hence v is even) and $v \neq 8$. There is a non-simple 5-polytope with v vertices and e edges, iff $\frac{1}{2}(5v + 3) \leq e \leq \binom{v}{2}$, and $(v, e) \neq (9, 25)$.

In particular, $\phi(v, 5) = \frac{1}{2}(5v + 3)$ if v is odd, and $\phi(v, 5) = 2.5v$ if v is even and not 8.

The values of $\phi(v, 6)$ are given by our previous results if $v \leq 14$. We can show that $\phi(v, 6) = 3v$ if $v = 15, 16, 17$ or if $v \geq 20$, and that $\phi(18, 6) = 56$ and $\phi(19, 6) = 59$. More precisely:

Theorem

There is a 6-polytope with v vertices and e edges, iff $\phi(v, 6) \leq e \leq \binom{v}{2}$, $e \neq 3v + 1$ and $(v, e) \neq (10, 34), (11, 36), (12, 38), (12, 39)$ or $(15, 47)$.

Finally, we mention a very recent result about impossible values of (v, e) .

Theorem

If $4 \leq k \leq d$, then a d -polytope with $v = d + k$ vertices cannot have between $\phi(v, d) + 1$ and $\phi(v, d) + k - 4$ edges.

In other words, if our polytope is not a multiplex, then it has at least $\max\{2, k - 3\}$ more edges than the multiplex.

